



 (02) 9625 2311
 (02) 9832 2469
 www.thevellagroup.com.au
 sales@thevellagroup.com.au
 PO Box 73, Doonside. NSW. 2767
 96 Glendenning Rd Plumpton. NSW. 2761

EQUINE MARE & FOAL

Typical Analysis (88% DM)	As Fed	Dry Matter
Digestible Energy	12.80	14.54
Protein (%)	18.00	20.45
Calcium (%)	1.70	1.70
Phosphorus (%)	0.84	0.95
Magnesium (%)	0.30	0.34
Sodium (%)	0.45	0.51
Chloride (%)	0.66	0.74
Added Salt (%)	1.00	
Lysine (%)	0.87	0.98
Potassium (%)	0.87	0.98
Fibre (%)	7.88	8.95
Fat (%)	6.20	7.04
Vit A IU/Kg	10,056	11,426
Vit D3 IU/Kg	2,500	2,840
Vit E mg/Kg	150.00	170.44
Vit K3 mg/Kg	1.25	1.42
Vit B1 mg/Kg	8.99	10.21

Vit B2 mg/Kg	7.72	8.77
Vit B6 mg/Kg	8.31	9.33
Vit B12 ug/Kg	12.50	14.20
Niacin mg/Kg	85.00	96.58
Biotin ug/Kg	268.00	304.52
Folic Acid mg/Kg	2.030	2.30
Iron mg/Kg	245.68	279.16
Zinc mg/Kg	114.68	130.31
Manganese mg/Kg	129.14	146.74
Copper mg/Kg	30.36	34.49
Cobalt mg/Kg	0.332	0.377
Selenium mg/Kg	0.655	0.744
Molybdenum mg/Kg	0.644	0.731
Iodine mg/Kg	0.840	0.954

Equine Mare & Foal has been formulated to suit the high demands of mares in the last trimester of pregnancy, lactating mares, foals, weanlings and yearlings.

Yea-Sacc™ Added to this diet

Other Specialty Diets

- Equine Cool - Active
- Equine Cool - Calm
- Livestock Maintenance Cubes

Category	Age	Weight	Kg's Equine Mare & Foal per day
Pregnant mares last trimester	-	550 kg	3.00 - 5.00
Lactating mares (0 - 3 months)	-	500 kg	3.00 - 5.00
Foals (creep feeding)	2 - 5 months	100 - 200 kg	0.50 - 1.50
Weanlings	5 - 12 months	200 - 300 kg	1.50 - 2.50
Yearlings	12 - 18 months	300 - 400 kg	2.50 - 3.50
Senior horses (maintenance - light work)	> 18 - 20 years	450 - 550 kg	2.50 - 3.50
Senior horses (light - moderate work)	> 18 - 20 years	450 - 550 kg	3.00 - 4.00
Increase in body condition	-	450 - 550 kg	3.00 - 4.00

- Senior horses can be fed low protein roughage due to high quality proteins in Mare & Foal. E.g. White chaffs and pasture hays preferred to lucerne and clover products.
- Feeding levels are intended as a guide only, ensure that clean, cool water and roughage in the form of pasture and/or hay is available at all times.
- Daily roughage intake should make up no less than 1.00% (minimum) - 1.50% (ideal) of the horses live body weight (LBW) to assist in maintaining healthy gut function.
- Pellets fed at levels greater than 1.00% of horses LBW should be fed over 2 or more feeds per day.
- Feeding levels based on other feeds available, body condition and rate of metabolism (good/poor doer) also on the maturity weight of the horse (500 - 550 kg).
- Introduce new feeds or changes to current feeding gradually over a 10 -14 day period.

Revision: 20070808