



(02) 9625 2311
 (02) 9832 2469
 www.thevellagroup.com.au
 sales@thevellagroup.com.au
 PO Box 73, Doonside. NSW. 2767
 96 Glendenning Rd Plumpton. NSW. 2761

EWE & LAMB

Typical Analysis	As Fed	Dry Matter
Energy MJ/ME kg	10.68	12.00
Protein (%)	16.00	18.00
Calcium (%)	1.78	2.00
Phosphorus (%)	0.53	0.60
Fibre (%)	4.27	4.80
Added Salt (%)	<1.00	

TRACE MINERAL PREMIX ADDED, FOR MORE DETAILED ANALYSIS PLEASE CONTACT VELLA STOCK FEEDS.

Ewe & Lamb has been formulated to assist in the growth of young lambs and the ewe to maintain condition whilst under heavy lactation. Also suitable for all sheep during times of drought or feeding off poor pastures to assist maintain condition or increase growth rates.

- Not suited to lambs under feedlot conditions, please see VSF Feedlot Lamb diet.

Other Specialty Products:

- Feedlot Lamb
- Cattle Fattener
- Livestock Maintenance Cube
- Beef Feedlot Concentrate
- McCoskers Brew
- Custom Diets

General Information

- Avoid feeding levels greater than 1.00 % of live body weight (LBW) - (1.50% max)
- Ensure roughage is available at all time especially when feeding pellet level is above 1.00 % LBW
- If unable to use self feeders, feed pellets on alternate days to avoid slug feeding
- Ewes in Lamb @ 0.75 - 1.50 kg per head per week
- Ewes after Lambing @ 0.75 - 1.50 kg per head per week
- Weaners @ 0.50 - 1.00 kg per head per day per week
- Rams in work @0.75 - 1.50 kg per head per week
- *Feeding levels are intended as a guide only, ensure that clean, cool water and roughage in the form of pasture and/or hay is available at all times.*
- *Results will vary according to seasonal conditions, age of animal, body condition prior to the introduction of supplements and other feed/roughage*
- *Daily roughage intake should make up no less that 1.50% of the livestock's, LBW (live body weight) to assist in maintaining healthy gut function and help avoid acidosis.*
- *For best results, feed should be introduced to livestock prior to other feeds becoming limited and/or animals start losing condition.*
- *Introduce new feeds or changes to current feeding gradually over a 14 - 21 day period.*
- *Phase out supplementary feeding gradually to help avoid set backs.*

Revision: 20080813