

## FREE RANGE LAYER

Typical Analysis	As Fed	Dry Matter
Energy KCAL / kg	2500	2770
Protein (min) (%)	15.50	17.20
Calcium (min) (%)	3.54	4.02
Phos (min) (%)	0.37	0.42
Fibre (max) (%)	6.00	6.60
Fat (max) (%)	6.00	6.60

TRACE MINERAL PREMIX ADDED, FOR MORE DETAILED ANALYSIS PLEASE CONTACT VELLA STOCK FEEDS.

**Free Range Layer** has been formulated to suit all breeds of layers from point of lay (18 - 20 weeks of age).

Energy density and analysis is well suited to young layers in early stages of lay, contains a yoke colour to assist in golden yokes.

NO antibiotics or growth promotants or hormones added.

- **Proteins** - No urea, all protein in the form of true proteins sources
- **Energy / Analysis** - For egg production and bird health
- **Trace Mineral Premix** - To assist in maintaining nutritional balance and layer health
- **Pellet** - easier acceptance and intake by layers, especially bantams
- **Clearly Identifiable Bags** - Avoids using the wrong product
- **Easy Lift Bag** - For your safety also in bulk and bulk bag form

### Other Specialty Products

- Chicken Starter
- Poultry Supreme
- Meat Bird Finisher
- Pullet Grower
- Free Range Scratch Mix
- Duck Grower
- Back Yard Layer
- Meat Bird Starter
- Specialty Diets

### General Information

- Introduce to birds from 18 - 20 weeks of age
- Intake varies from 95 grams - 110 grams per day depending on bird health & breed, lighting, housing, climate conditions and water supply
- Feeding hens to production is important to maximise bird health, egg production & yoke colour
- Fresh, clean water to be accessible at all times
- Greens can be fed in conjunction with pellets
- Older hens (>2 years) egg production will decline rapidly due to aging

- *Egg production rates will vary according to a range of external factors including bird age, housing, lighting, variable weather conditions, predators, inoculations and hygiene to name a few*
- *Phase supplementary feeding both in and out gradually to help avoid set backs.*
- *Housing should be well vented but draft free*
- *Ensure clean, cool water is accessible at all times*
- *Feeders and waterers need to be easily accessible and cleaned of litter on a regular basis*
- *Quarantine sick layers to prevent possible outbreaks*
- *Dead layers need to be removed straight away and disposed of.*

Revision: 20170906