

MEAT BIRD FINISHER - BREEDER

Typical Analysis	As Fed	Dry Matter
Energy KCAL / kg	2780	3080
Protein (min) (%)	19.00	21.10
Calcium (min) (%)	1.67	1.90
Phos (min) (%)	0.88	1.00
Fibre (max) (%)	6.50	7.20
Fat (max) (%)	7.00	7.70

TRACE MINERAL PREMIX ADDED. FOR MORE DETAILED ANALYSIS PLEASE CONTACT VELLA STOCK FEEDS.

Meat Bird Grower / Finisher has been formulated to suit

Turkeys (7 - 8 weeks to desired weight)

Broilers (4 - 5 weeks to desired weight)

Game Birds (Quail, Guinea Fowl & Pheasants, 4 - 5 weeks to desired weight)

- **Probiotic and Prebiotic** - Good bacteria for gut health
- **Coccidiostat** - Regano, a natural herbal coccidiostat used, no withholding period for meat and or egg consumption. Also suitable for ducklings to consume.
- **Natural Betaine** - Support chicks during times of heat stress
- **Vegetable Proteins** - No restricted animal materials (RAM) used in our diets
- **Vitamins** - Species specific vitamins & minerals included to support health and production
- **Antioxidant** - Assist health
- **No** hormones or antibiotics used

Other Specialty Products

- Chick Starter
- Free Range Layer
- Meat Bird Starter
- Pullet Grower
- Poultry Supreme
- Duck Grower
- Back Yard Layer
- Free Range Scratch Mix
- Specialty Diets

General Information

- If you desire commercial growth rates we suggest that additional Calcium be made available to suit individual bird requirements
- The diet should be fed ad-lib to all birds 24 / 7 with adequate feeder space for stock
- A natural herbal (Regano) coccidiostat has been added to assist in the control of Coccidiosis.
- Fresh, clean water to be accessible at all times
- Remove unused wasted feed on a regular basis
- Feed intakes will vary at times, due to climate, bird health / stress and general housing and access to feeders
- Meat Bird is designed for the domestic grower, individual specie commercial diets available in bulk

- *Growth rates will vary according to a range of external factors including housing, variable weather conditions, predators, inoculations and hygiene to name a few*
- *Phase supplementary feeding both in and out gradually to help avoid set backs.*
- *Young stock are very sceptical to cold and draughts, housing should be well vented but draught free*
- *Ensure clean, cool water is accessible at all times*
- *Feeders and waterers needs to be easily accessible and cleaned of litter on a regular basis*
- *Quarantine sick birds to prevent possible outbreaks*

Revision: 20170906