



(02) 9625 2311
 (02) 9832 2469
www.thevellagroup.com.au
sales@thevellagroup.com.au
 PO Box 73, Doonside. NSW. 2767
 96 Glendenning Rd Plumpton. NSW. 2761

PULLET GROWER

Typical Analysis	As Fed	Dry Matter
Energy KCAL / kg	2600	2880
Protein (min) (%)	16.00	17.70
Calcium (min) (%)	1.31	1.9
Phos (min) (%)	0.48	0.55
Fibre (max) (%)	6.00	6.60
Fat (max) (%)	6.00	6.60

TRACE MINERAL PREMIX ADDED, FOR MORE DETAILED ANALYSIS PLEASE CONTACT VELLA STOCK FEEDS.

Pullet Grower has been formulated to suit all breeds of pullets from 6 - 7 weeks to point of lay (18 - 20 weeks of age). This feed contains a natural herbal coccidiostat.

This is a complete feed and no other supplement is required

- **Probiotic and Prebiotic** - Good bacteria for gut health
- **Coccidiostat** - Regano, a natural herbal coccidiostat used, no withholding period for meat and or egg consumption. Also suitable for ducklings to consume.
- **Natural Betaine** - Support chicks during times of heat stress
- **Vegetable Proteins** - No restricted animal materials (RAM) used in our diets
- **Vitamins** - Species specific vitamins & minerals included to support health and production
- **Antioxidant** - Assist health
- **No** added hormones or antibiotics used

Other Specialty Products

- Chicken Starter
- Poultry Supreme
- Meat Bird Finisher / Breeder
- Back Yard Layer
- Free Range Scratch Mix
- Duck Grower
- Free Range Layer
- Meat Bird Starter
- Specialty Diets

General Information

- A complete diet available in a course crumble form, providing important nutrients for growing pullets
- Suitable to be fed to all breeds of birds from 6 weeks to point of lay (18—20 weeks)
- A coccidiostat has been added to assist in the control of Coccidiosis
- Fresh, clean water to be accessible at all times
- Feed intake will vary at times, due to climate and general housing of pullets
- As pullets develop (age) feed intake will increase rapidly until maturity
- Average consumption from 6 weeks of age to point of lay, approximately 5-7 kg of feed per bird

- *Growth rates will vary according to a range of external factors including housing, variable weather conditions, predators, inoculations and hygiene to name a few*
- *Phase supplementary feeding both in and out gradually to help avoid set backs.*
- *Housing should be well vented but draft free*
- *Ensure clean, cool water is accessible at all times*
- *Feeders and waterers needs to be easily accessible and cleaned of litter on a regular basis*
- *Quarantine sick pullets to prevent possible outbreaks*
- *Dead pullets need to be removed straight away and disposed of.*

Revision: 20170906