



96 Glendenning Rd Plumpton NSW 2761  
 PO Box 73 Doonside NSW 2767  
 02 9625 2311  
 info@thevellagroup.com.au  
 www.thevellagroup.com.au  
 www.facebook.com/vellastockfeeds

## EWE & LAMB GROWER - BREEDER

Typical Analysis	As Fed	Dry Matter
Energy MJ/ME kg	10.80	12.00
Protein (%)	16.20	18.00
Calcium (%)	1.30	1.44
Phosphorus (%)	0.65	0.72
Fibre Max (%)	10.00	11.11
Sodium Max (%)	2.50	

*TRACE MINERAL PREMIX ADDED, FOR MORE DETAILED ANALYSIS PLEASE CONTACT VELLA STOCK FEEDS.*

**Ewe & Lamb** has been formulated to assist in the growth of young lambs and the ewe to maintain condition whilst under heavy lactation. Also suitable for all sheep during times of drought or feeding off poor pastures to assist maintain condition or increase growth rates.

- **True Proteins** - Proteins from natural sources
- **No Urea** - Reduces possible conflict if feeding Urea blocks / supplements with paddock roughage
- **Energy** - To assist with growth and maintenance of the ewe whilst lactating
- **Trace Mineral Premix** - To assist in maintaining correct nutrition balance.
- **Pellet** - 4.2 mm diameter pellet, less wastage and can be trail fed.
- **Packaging** - Bags, Bulk Bags & Bulk Bulk & Bulk Bags minimum 3 tonne order

### Typical raw materials considered in the formulation of this diet

Cereal Grains & their Co-Products, Lupins, Canola & Soybean Meal, Vegetable Oils, Amino Acids, Calcium, Phosphorus, Vitamin Premix, Salt, Sodium Bicarbonate, Sodium Bentonite

All raw materials used are steamed cooked prior to the pelleting process

### General Information

- Avoid feeding levels greater than 1.00 % of live body weight (LBW) - (1.50% max)
  - Ensure roughage is available at all time especially when feeding pellet level is above 1.00 % LBW
  - If unable to use self feeders, feed pellets on alternate days to avoid slug feeding
  - Ewes in Lamb @ 0.75 - 1.50 kg per head per week
  - Ewes after Lambing @ 0.75 - 1.50 kg per head per week
  - Weaners @ 0.50 - 1.00 kg per head per day per week
  - Rams in work @0.75 - 1.50 kg per head per week
- 
- *Feeding levels are intended as a guide only, ensure that clean, cool water and roughage in the form of pasture and/or hay is available at all times.*
  - *Results will vary according to seasonal conditions, age of animal, body condition prior to the introduction of supplements and other feed/roughage*
  - *Daily roughage intake should make up no less that 1.50% of the livestock's, LBW (live body weight) to assist in maintaining healthy gut function and help avoid acidosis.*
  - *For best results, feed should be introduced to livestock prior to other feeds becoming limited and/or animals start losing condition.*
  - *Introduce new feeds or changes to current feeding gradually over a 14 - 21 day period.*
  - *Phase out supplementary feeding gradually to help avoid set backs.* Revision: 20240404