

96 Glendenning Rd Plumpton NSW 2761
 PO Box 73 Doonside NSW 2767
 02 9625 2311

info@thevellagroup.com.au
www.thevellagroup.com.au

www.facebook.com/vellastockfeeds

CATTLE SUPREME GROWER-BREEDER

| Typical Analysis | As Fed | Dry Matter |
|----------------------|--------|------------|
| Energy MJ/kg (min) | 11.25 | 12.50 |
| Protein (%) (min) | 16.00 | 17.75 |
| Calcium (%) (min) | 1.20 | 1.30 |
| Phosphorus (%) (min) | 0.60 | 0.66 |
| NDF (%) (max) | 27.00 | 30.00 |
| Added Salt (%) (max) | 1.70 | |

<u>Cattle Supreme</u> has been specifically formulated for the use on young, growing, showing and fattening (finish off) cattle. Also well suited to lactating cattle, stud cattle being prepped for show or sales. Can also be used for bulls both working or maintenance.

- Proteins No urea, added protein in the form quality vegetable proteins sources
- Energy Multiple energy sources used to complement protein sources to further assist with sustainable growth rates and bloom
- Trace Mineral Premix To assist in maintaining nutritional balance, livestock health and coat condition
- Buffers Additional buffers, Sodium Bicarbonate & Acid Buf added to assist in maintaining rumen function

Typical raw materials considered in the formulation of this diet

Wheat, Barley, Sorghum & their Co-Products, Lupins, Canola & Soybean Meal, Vegetable Oils, Calcium, Phosphorus, Vitamin Premix, Salt, Sodium Bicarbonate, Acid Buf, Sodium Bentonite, Antioxidant, Molasses All raw materials used are steamed cooked prior to the pelleting precess

| Lactating Cows | To assist the dam with milk supply and lessen potential weight loss from the demand of the calf, feed pellets at 1 - 4 kg per head per day, not to exceed 1% of LBW (§) | |
|--------------------------|---|--|
| Weaners | To avoid potential "set backs" at time of weaning feed pellets at 0.5 - 1% of LBW (§) | |
| Growing Cattle | Pending desired growth rates required, feed pellets to a maximum of 1% LBW (§) | |
| Show & Sale of Cattle | To assist with top line condition and bloom, feed pellets at 2 - 4 kg per head per day, starting a minimum of three weeks prior to show, ensuring, not to exceed 1% of their LBW (§). Oil seeds may be used to complement pellets to further enhance coat condition, avoid feeding levels of oil seeds greater than 250 grams per head per day. | |
| Fattening Cattle | Please contact Vella Stock Feeds for individual feeding suggestions best suited to your requirements for the fattening / finishing off, of cattle | |
| Bulls | Working bulls to be feed to level of activity, generally, bulls in high work to be fed pellets at a maximum of 0.5% LBW or 4 kg whichever is the lesser amount (§) | |

(§) Contact Vella stock Feeds for a feeding program if intending to feed levels greater than 1.00% of Live Body Weight (LBW) per 24 hour period e.g. 300 kg beast x 1% = 3.00 kg

- Access to clean, palatable roughage (24/7). Roughage to be at least 25 mm (1") or longer (effective fibre).
- Cattle should have access to at least 2% of their LBW of roughage daily (24/7)
- Cool, clean and accessible water to be available at all times (24/7)
- Introduce feed in a step up program, increasing feed levels to desired level (§) over a 14 21 day period
- Keep yards and feeding equipment / area hygienic by removing feed and animal waste regularly
- Monitor cattle's dung, if dung becomes very loose this <u>MAY</u> be an indication of cattle over consuming pelleted feed