



 (02) 9625 2311
 (02) 9832 2469
 www.thevellagroup.com.au
 sales@thevellagroup.com.au
 PO Box 73, Doonside. NSW. 2767
 96 Glendenning Rd Plumpton. NSW. 2761

ALPACA PELLETS

Typical Analysis	As Fed	Dry Matter
Energy MJ/ME kg	11.00	12.20
Protein (Min) (%)	16.00	17.70
Calcium (Min) (%)	1.50	1.68
Phos (Min) (%)	0.80	0.90
Fibre (Max) (%)	10.00	10.90
Added Salt (%)	< 1.50	

TRACE MINERAL PREMIX ADDED. FOR MORE DETAILED ANALYSIS PLEASE CONTACT VELLA STOCK FEEDS.

Alpaca Pellets have been formulated as a supplement to be fed in conjunction with pasture and/or hay to young, growing and mature stock.

Also suitable to be fed to Llamas

Typical raw materials considered in the formulation of this diet

Wheat, Barley, Sorghum & their Co-Products, Lupins, Canola & Soybean Meal, Lucerne Meal, Vegetable Oils, Amino Acids, Probiotics, Calcium, Phosphorus, Vitamin Premix, Salt, Sodium Bicarbonate, Sodium Bentonite, Potassium Chloride, Antioxidant

All raw materials used are steamed cooked prior to the pelleting process

Feeding Suggestions

- Estimated dry matter intake (kg) of Alpacas and Llamas is approximately 1.80% - 2.00% of LBW (live body weight) per day
- Pelleted feed should not exceed 50% (dry matter) of total intake
- Where possible feed allocation over two or more feeds per day
- Where it is necessary to feed mature Alpacas (>0.50 kg) and Llamas (>1.00 kg) per day, mix equal amounts (by weight) of a course roughage (e.g. white chaff) with pellets
- Ensure that stock have access to suitable (palatable) roughage in the form of pasture and or hay at all times
- Access to cool, clean drinking water at all times
- Introduce pellets slowly and remove any waste daily
- *Feeding levels are intended as a guide only, ensure that clean, cool water and roughage in the form of pasture and/or hay is available at all times.*
- *Results will vary according to seasonal conditions, age of animal, housing, health and body condition prior to the introduction of supplements and other feed / roughage*
- *For best practice, feed should be introduced to livestock prior to other feeds becoming limited*
- *Phase new feeds in and out gradually over a 14 - 21 day period to avoid set backs.*

Revision; 20190517