

EQUINE COOL - CALM

Typical Minimum Analysis	Dry Matter
Digestible Energy	12.50
Protein (%)	14.00
Calcium (%)	1.70
Phosphorus (%)	0.83
Magnesium (%)	0.40
Sodium (%)	0.50
Chloride (%)	0.70
Added Salt (%) (Max)	<1.00%
Lysine (%)	0.60
Potassium (%)	0.90
Fibre (%) (Max)	12.00
Fat (%) (Max)	5.00
Vit A IU/Kg	3340
Vit D3 IU/Kg	830
Vit E mg/Kg	33
Vit B2 mg/Kg	5.50
Vit B6 mg/Kg	8.70
Vit B12 ug/Kg	4.00

Niacin mg/Kg	105.00
Biotin mg/Kg	0.25
Folic Acid mg/Kg	0.45
Iron mg/Kg	264.00
Zinc mg/Kg	105.00
Manganese mg/Kg	130.00
Copper mg/Kg	15.50
Cobalt mg/Kg	0.20
Selenium mg/Kg	0.65
Molybdenum mg/Kg	0.55
Iodine mg/Kg	0.25

Equine Cool Calm

Has been specifically formulated to suit the needs of all mature ponies and horses from maintenance to medium work loads e.g. pony club, hacking, trail riding, dressage. Correct equine nutrition will assist in maintaining, improving wellbeing, tone and coat condition, giving you the upper hand in today's competitive competitions. Nutritional support is available for users of any of our specialty equine diets.

Typical raw materials considered in the formulation of this diet

Steamed Barley, Bran & Pollard, Canola, Vegetable Oils, Amino Acids, Calcium, Phosphorus, Species Specific Vitamin Premix, Salt, Sodium Bentonite, Antioxidant, Molasses
 All raw materials used are steamed cooked prior to the pelleting process

Estimated Weight	Maintenance—Light Work	Light—Moderate Work
200 kg	0.50 - 1.50 kg	0.90 - 2.00 kg
250 kg	0.85 - 2.00 kg	1.25 - 2.50 kg
350 kg	1.20 - 2.80 kg	1.75 - 3.50 kg
450 kg	1.50 - 3.60 kg	2.25 - 4.25 kg
550 kg	2.00 - 4.50 kg	2.75 - 4.75 kg

- Suggested feeding levels are intended as a guide only, ensure adequate clean, cool water and roughage as pasture and or hay is available at all times.
- Daily roughage intake should make up no less than 1.0% (minimum) -1.50% (ideal) of pony/horse, live body weight to assist in maintaining healthy gut function.
- Pellets fed at levels equal to or greater than 1% of pony/horses live body weight should be fed over two or more feeds per day.
- Feeding levels dependent on the availability of other feeds, body condition and rate of metabolism
- Introduce new feeds or changes gradually over a 10 -14 day period.
- Free feeding program service available to assist with individual needs and goals, please contact mill for details.