

MEAT BIRD STARTER

Typical Analysis	As Fed	Dry Matter
Energy KCAL / kg	2900	3220
Protein (min) (%)	23.00	25.50
Calcium (min) (%)	1.00	1.10
Phos (min) (%)	0.80	0.88
Fibre (max) (%)	6.00	6.60
Fat (max) (%)	5.00	5.50

TRACE MINERAL PREMIX ADDED, FOR MORE DETAILED ANALYSIS PLEASE CONTACT VELLA STOCK FEEDS.

Meat Bird Starter has been formulated to suit

Poults (day old to 6 - 8 weeks)

Broilers (day old to 4 - 5 weeks)

Game Birds (Quail, Guinea Fowl & Pheasants, day old to 4 - 6 weeks of age)

- **Probiotic and Prebiotic** - Good bacteria for gut health
- **Coccidiostat** - Regano, a natural herbal coccidiostat used, no withholding period for meat and or egg consumption. Also suitable for ducklings to consume.
- **Natural Betaine** - Support chicks during times of heat stress
- **Vegetable Proteins** - No restricted animal materials (RAM) used in our diets
- **Vitamins** - Species specific vitamins & minerals included to support health and production
- **Antioxidant** - Assist health
- **No** added hormones or antibiotics used

Typical raw materials considered in the formulation of this diet

Wheat, Barley, Sorghum & their Co-Products, Lupins, Corn, Canola & Soybean Meal, Lucerne Meal, Vegetable Oils, Amino Acids, Probiotics, Prebiotics, Calcium, Phosphorus, Species Specific Vitamin Premix, Salt, Sodium Bicarbonate, Sodium Bentonite, Natural Betaine, Antioxidant, Regano (Herbal Coccidiostat), Natural Digestive Enzymes, Molasses
 All raw materials used are steamed cooked prior to the pelleting process

General Information

- A complete commercial crumble (diet) providing all requirements for young growing stock
- The diet should be fed ad-lib to all birds
- Fresh, clean water to be accessible at all times and where possible the supplementation of the water with a water soluble vitamins in the first 7 days of life will aid less dominant stock.
- Assist young stock in the first few days of life, change feed regularly
- Feed intakes will vary at times, due to climate, bird health / stress and general housing and access to feeding and watering stations.
- The sprinkling of the starter diet on paper in the brooder for the first week will also aid less dominate stock.

- *Growth rates will vary according to a range of external factors including housing, variable weather conditions, predators, inoculations and hygiene to name a few*
- *Phase supplementary feeding both in and out gradually to help avoid set backs.*
- *Young stock are very sceptical to cold and draughts, housing should be well vented but draught free*
- *Ensure clean, cool water is accessible at all times*
- *Feeders and waterers needs to be easily accessible and cleaned of litter on a regular basis*
- *Quarantine sick birds to prevent possible outbreaks*

Revision: 20200303